



## Pediatric COVID-19 Vaccine Information (Children 5 Years of Age and Older)

Children 5 Years of Age and Older Updated (2023–2024 Formula) Pfizer-BioNTech COVID-19 Vaccine for those who are NOT moderately or severely immunocompromised\*

COVID-19 vaccination history <sup>†</sup> (regardless of COVID-19 vaccine formula)	Schedule for administration of 2023-24 Pfizer-BioNTech COVID-19 Vaccine
Unvaccinated	Give 1 dose now.
Any number of previous doses of COVID-19 vaccine, <b>NOT</b> including at least 1 dose of 2023–24 COVID-19 vaccine	Give 1 dose at least 8 weeks (2 months) after the previous dose.
Any number of previous doses COVID-19 vaccine, <b>INCLUDING</b> at least 1 dose of 2023–24 COVID-19 vaccine	No further doses are indicated.

\* Persons with a recent SARS-CoV-2 infection may consider delaying vaccination by 3 months from symptom onset or positive test (if infection was asymptomatic).

† COVID-19 vaccination history refers to previous receipt of dose(s) of Original monovalent mRNA, bivalent mRNA vaccine, Updated (2023–2024 Formula), or a combination of the three, unless otherwise specified.

**Children 5 Years of Age and Older Updated (2023–2024 Formula) Pfizer-BioNTech COVID-19 Vaccine for those who ARE moderately or severely immunocompromised\***

COVID-19 vaccination history* (regardless of COVID-19 vaccine formula)	Schedule for administration of 2023-24 Pfizer-BioNTech COVID-19 Vaccine
Unvaccinated	Give a 3-dose initial series. Administer: <ul style="list-style-type: none"> <li>▪ Dose 1 now</li> <li>▪ Dose 2 at least 3 weeks after Dose 1</li> <li>▪ Dose 3 at least 4 weeks after Dose 2</li> </ul>
1 previous dose of any Pfizer-BioNTech COVID-19 Vaccine (Dose 1) <sup>†</sup>	Complete series. Administer: <ul style="list-style-type: none"> <li>▪ Dose 2 at least 3 weeks after Dose 1</li> <li>▪ Dose 3 at least 4 weeks after Dose 2</li> </ul>
2 doses of any Pfizer-BioNTech COVID-19 Vaccine (Doses 1 and 2) <sup>†</sup>	Complete series. Administer: <ul style="list-style-type: none"> <li>▪ Dose 3 at least 4 weeks after Dose 2</li> </ul>
3 or more doses of Pfizer-BioNTech COVID-19 Vaccine, <b>NOT</b> including at least 1 dose of 2023–24 COVID-19 vaccine <sup>†</sup>	Give 1 dose at least 8 weeks (2 months) after the previous dose.
3 or more doses of Pfizer-BioNTech COVID-19 Vaccine, <b>INCLUDING</b> at least 1 dose of 2023–24 COVID-19 vaccine <sup>†</sup>	<ul style="list-style-type: none"> <li>▪ People who are moderately or severely immunocompromised have the option to receive 1 additional dose at least 8 weeks (2 months) following the last recommended dose.</li> <li>▪ Further additional dose(s) may be administered, informed by the clinical judgement of a health care provider and personal preference and circumstances.</li> <li>▪ Any further additional doses should be administered at least 8 weeks (2 months) after the last COVID-19 vaccine dose.</li> </ul>

\* COVID-19 vaccination history refers to previous receipt of dose(s) of Original monovalent mRNA, bivalent mRNA vaccine, Updated (2023–2024 Formula), or a combination of the three, unless otherwise specified.

† People who are recommended to receive a multidose mRNA series for initial vaccination (i.e., children ages 6 months–4 years and people who are moderately or severely immunocompromised) should receive all doses from the same manufacturer. However, in the following exceptional situations a different age-appropriate COVID-19 vaccine product may be administered: the same vaccine is not available, the person would otherwise not complete the vaccination series, or the person starts but is unable to complete a vaccination series with the same vaccine due to a contraindication.